

# LARGE Publications

2016

Nakao H, Harada T, Nakao K, Kiyonari H, Inoue K, Furuta Y, Aiba A. (2016) A possible aid in targeted insertion of large DNA elements by CRISPR/Cas in mouse zygotes. *Genesis*. 54(2):65-77.

Shi D, Usami F, Komatsu K, Oka S, Abe T, Uemura T, Fujimori T. (2016) Dynamics of planar cell polarity protein Vangl2 in the mouse oviduct epithelium. *Mech Dev*. 141:78-89.

Ibeh BO, Furuta Y, Habu JB, Ogbadu L. (2016) Humanized mouse as an appropriate model for accelerated global HIV research and vaccine development: current trend. *Immunopharmacol Immunotoxicol*. 1-13.

Toya, M., Kobayashi, S., Kawasaki, M., Shioi, G., Kaneko, M., Ishiuchi, T., Misaki, K., Meng, W, and Takeichi, M. (2016) CAMSAP3 orients the apical-to-basal polarity of microtubule arrays in epithelial cells. *Proc Natl Acad Sci U S A*. 113(2):332-7.

Mukai, H., Muramatsu, A., Mashud, R., Kubouchi, K., Tsujimoto, S., Hongu, T., Kanaho, Y., Tsubaki, M., Nishida, S., Shioi, G., Danno, S., Mehruba, M., Satoh, R. and Sugiura, R. (2016) PKN3 is the major regulator of angiogenesis and tumor metastasis in mice. *Sci Rep*. 6:18979. doi: 10.1038/srep18979.

Watanabe S, Shioi G, Furuta Y, Goshima G. (2016) Intra-spindle Microtubule Assembly Regulates Clustering of Microtubule-Organizing Centers during Early Mouse Development. *Cell Rep*. 15(1):54-60.

Isozaki T, Komenoi S, Lu Q, Usuki T, Tomokata S, Matsutomo D, Sakai H, Bando K, Kiyonari H, Sakane F. (2016) Deficiency of diacylglycerol kinase  $\eta$  induces lithium-sensitive mania-like behavior. *J Neurochem*. 138(3), 448-456.

Ip JY, Sone M, Nashiki C, Pan Q, Kitaichi K, Yanaka K, Abe T, Takao K, Miyakawa T, Blencowe BJ, Nakagawa S. (2016) Gomafu lncRNA knockout mice exhibit mild hyperactivity with enhanced responsiveness to the psychostimulant methamphetamine. *Sci Rep.* 6:27204. doi: 10.1038/srep27204.

Hata E, Sasaki N, Takeda A, Tohya K, Umemoto E, Akahoshi N, Ishii S, Bando K, Abe T, Kano K, Aoki J, Hayasaka H, Miyasaka M. (2016) Lysophosphatidic acid receptors LPA4 and LPA6 differentially promote lymphocyte transmigration across high endothelial venules in lymph nodes. *Int Immunol.* 28(6):283-92.

Haraguchi R, Kitazawa R, Mori K, Tachibana R, Kiyonari H, Imai Y, Abe T, Kitazawa S. sFRP4-dependent Wnt signal modulation is critical for bone remodeling during postnatal development and age-related bone loss. *Sci Rep.* 2016 Apr 27;6:25198. doi: 10.1038/srep25198.

Tokumasu R, Yamaga K, Yamazaki Y, Murota H, Suzuki K, Tamura A, Bando K, Furuta Y, Katayama I, Tsukita S. (2016) Dose-dependent role of claudin-1 in vivo in orchestrating features of atopic dermatitis. *Proc Natl Acad Sci U S A.* 113(28):E4061-8. doi: 10.1073/pnas.1525474113.

Hata K, Yanase N, Sudo K, Kiyonari H, Mukumoto Y, Mizuguchi J, Yokosuka T. (2016) Differential regulation of T-cell dependent and T-cell independent antibody responses through arginine methyltransferase PRMT1 in vivo. *FEBS Lett.* 590(8):1200-10. doi: 10.1002/1873-3468.12161.

Doi M, Murai I, Kunisue S, Setsu G, Uchio N, Tanaka R, Kobayashi S, Shimatani H, Hayashi H, Chao HW, Nakagawa Y, Takahashi Y, Hotta Y, Yasunaga J, Matsuoka M, Hastings MH, Kiyonari H, Okamura H. (2016) Gpr176 is a Gz-linked orphan G-protein-coupled receptor that sets the pace of circadian behaviour. *Nat Commun.* 7:10583. doi: 10.1038/ncomms10583.

Nishi K, Sato Y, Ohno M, Hiraoka Y, Saijo S, Sakamoto J, Chen PM, Morita Y, Matsuda S, Iwasaki K, Sugizaki K, Harada N, Mukumoto Y, Kiyonari H, Furuyama K,

Kawaguchi Y, Uemoto S, Kita T, Inagaki N, Kimura T, Nishi E. (2016) Nardilysin Is Required for Maintaining Pancreatic  $\beta$ -Cell Function. *Diabetes*. 65(10):3015-27.

Connor SA, Ammendrup-Johnsen I, Chan AW, Kishimoto Y, Murayama C, Kurihara N, Tada A, Ge Y, Lu H, Yan R, LeDue JM, Matsumoto H, Kiyonari H, Kirino Y, Matsuzaki F, Suzuki T, Murphy TH, Wang YT, Yamamoto T, Craig AM. (2016) Altered Cortical Dynamics and Cognitive Function upon Haploinsufficiency of the Autism-Linked Excitatory Synaptic Suppressor MDGA2. *Neuron*. 91(5):1052-68.

Usuki T, Takato T, Lu Q, Sakai H, Bando K, Kiyonari H, Sakane, F. (2016) Behavioral and pharmacological phenotypes of brain-specific diacylglycerol kinase  $\delta$ -knockout mice. *Brain Res*. 1648 (Part A), 193–201.

Makihara H, Nakai S, Ohkubo W, Yamashita N, Nakamura F, Kiyonari H, Shioi G, Jitsuki-Takahashi A, Nakamura H, Tanaka F, Akase T, Kolattukudy P, Goshima Y. (2016) CRMP1 and CRMP2 have synergistic but distinct roles in dendritic development. *Genes Cells*. 21(9):994-1005.

Miyake K, Ohta T, Nakayama H, Doe N, Terao Y, Oiki E, Nagatomo I, Yamashita Y, Abe T, Nishikura K, Kumanogoh A, Hashimoto K, Kawahara Y. (2016) CAPS1 RNA Editing Promotes Dense Core Vesicle Exocytosis. *Cell Reports* 17(8): 2004-2014.

Higuchi M, Kawamura H, Matsuki H, Hara T, Takahashi M, Saito S, Saito K, Jiang S, Naito M, Kiyonari H, Fujii M. (2016) USP10 Is an Essential Deubiquitinase for Hematopoiesis and Inhibits Apoptosis of Long-Term Hematopoietic Stem Cells. *Stem Cell Reports*. 7(6):1116-1129.

Kusano R, Fujita K, Shinoda Y, Nagaura Y, Kiyonari H, Abe T, Watanabe T, Matsui Y, Fukaya M, Sakagami H, Sato T, Funahashi JI, Ohnishi M, Tamura S, Kobayashi T. (2016) Targeted disruption of the mouse protein phosphatase *ppm1l* gene leads to structural abnormalities in the brain. *FEBS Lett*. 590(20):3606-3615.

## 2015

Hoshino, H., Shioi, G. and Aizawa, S. (2015) AVE protein expression and visceral endoderm cell behavior during anterior-posterior axis formation in mouse embryos: Asymmetry in OTX2 and DKK1 expression. *Dev Biol* 402(2): 175-91.

Hiramuki, Y., Sato, T., Furuta, Y., Surani, M. A. and Sehara-Fujisawa, A. (2015) Mest but Not MiR-335 Affects Skeletal Muscle Growth and Regeneration. *PLoS One* 10(6): e0130436.

Oshiro, H., Hirabayashi, Y., Furuta, Y., Okabe, S. and Gotoh, Y. (2015) Up-regulation of HP1gamma expression during neuronal maturation promotes axonal and dendritic development in mouse embryonic neocortex. *Genes Cells* 20(2): 108-20.

Sato, S., Yajima, H., Furuta, Y., Ikeda, K. and Kawakami, K. (2015) Activation of Six1 Expression in Vertebrate Sensory Neurons. *PLoS One* 10(8): e0136666.

Fukuda, N., Naito, S., Masukawa, D., Kaneda, M., Miyamoto, H., Abe, T., Yamashita, Y., Endo, I., Nakamura, F. and Goshima, Y. (2015) Expression of ocular albinism 1 (OA1), 3, 4- dihydroxy- L-phenylalanine (DOPA) receptor, in both neuronal and non-neuronal organs. *Brain Res* 1602: 62-74.

Recuenco, M. C., Ohmori, T., Tanigawa, S., Taguchi, A., Fujimura, S., Conti, M. A., Wei, Q., Kiyonari, H., Abe, T., Adelstein, R. S. and Nishinakamura, R. (2015) Nonmuscle Myosin II Regulates the Morphogenesis of Metanephric Mesenchyme-Derived Immature Nephrons. *J Am Soc Nephrol* 26(5): 1081-91.

Tsujimura, K., Irie, K., Nakashima, H., Egashira, Y., Fukao, Y., Fujiwara, M., Itoh, M., Uesaka, M., Imamura, T., Nakahata, Y., Yamashita, Y., Abe, T., Takamori, S. and Nakashima, K. (2015) miR-199a Links MeCP2 with mTOR Signaling and Its Dysregulation Leads to Rett Syndrome Phenotypes. *Cell Rep* 12(11): 1887-901.

Arimoto, K., Hishiki, T., Kiyonari, H., Abe, T., Cheng, C., Yan, M., Fan, J. B.,

Futakuchi, M., Tsuda, H., Murakami, Y., Suzuki, H., Zhang, D.E. and Shimotohno, K. (2015) Murine Herc6 Plays a Critical Role in Protein ISGylation In Vivo and Has an ISGylation-Independent Function in Seminal Vesicles. *J Interferon Cytokine Res* 35(5): 351-8.

Tanaka, H., Takechi, M., Kiyonari, H., Shioi, G., Tamura, A. and Tsukita, S. (2015) Intestinal deletion of Claudin-7 enhances paracellular organic solute flux and initiates colonic inflammation in mice. *Gut* 64(10): 1529-38.

Nojiri, T., Hosoda, H., Tokudome, T., Miura, K., Ishikane, S., Otani, K., Kishimoto, I., Shintani, Y., Inoue, M., Kimura, T. Sawabata, N., Minami, M., Nakagiri, T., Funaki, S., Takeuchi, Y., Maeda, H., Kidoya, H., Kiyonari, H., Shioi, G., Arai, Y., Hasegawa, T., Takakura, N., Hori, M., Ohno, Y., Miyazato, M., Mochizuki, N., Okumura, M. and Kangawa, K. (2015) Atrial natriuretic peptide prevents cancer metastasis through vascular endothelial cells. *Proc Natl Acad Sci U S A* 112(13): 4086-91.

Lee, M. S., Hwang, K. S., Oh, H. W., Ji-Ae, K., Kim, H. T., Cho, H. S., Lee, J. J., Yeong Ko, J., Choi, J. H., Jeong, Y. M., You, K.H., Kim, J., Park, D.S., Nam, K.H., Aizawa, S., Kiyonari, H., Shioi, G., Park, J.H., Zhou, W., Kim, N.S. and Kim, C.H. (2015) IFT46 plays an essential role in cilia development. *Dev Biol* 400(2): 248-57.

Imai, Y., Kobayashi, Y., Inoshita, T., Meng, H., Arano, T., Uemura, K., Asano, T., Yoshimi, K., Zhang, C. L., Matsumoto, G., Ohtsuka, T., Kageyama, R., Kiyonari, H., Shioi, G., Nukina, N., Hattori, N. and Takahashi, R. (2015) The Parkinson's Disease-Associated Protein Kinase LRRK2 Modulates Notch Signaling through the Endosomal Pathway. *PLoS Genet* 11(9): e1005503.

Imai, N., Suzuki, M., Hayashi, K., Ishigami, M., Hirooka, Y., Abe, T., Shioi, G., Goto, H. and Fujimoto, T. (2015) Hepatocyte-Specific Depletion of UBXD8 Induces Periportal Steatosis in Mice Fed a High-Fat Diet. *PLoS One* 10(5): e0127114.

Nakatsu, Y., Iwashita, M., Sakoda, H., Ono, H., Nagata, K., Matsunaga, Y., Fukushima, T., Fujishiro, M., Kushiyama, A., Kamata, H., Takahashi, SI., Katagiri, H., Honda, H.,

Kiyonari, H., Uchida, T. and Asano, T. (2015) Prolyl Isomerase Pin1 Negatively Regulates AMPK by Associating with the CBS Domain in the gamma-subunit. *J Biol Chem.* 290(40):24255-66.

Kurimoto, K., Yabuta, Y., Hayashi, K., Ohta, H., Kiyonari, H., Mitani, T., Moritoki, Y., Kohri, K., Kimura, H., Yamamoto, T., Katou, Y., Shirahige, K. and Saitou, M. (2015) Quantitative Dynamics of Chromatin Remodeling during Germ Cell Specification from Mouse Embryonic Stem Cells. *Cell Stem Cell* 16(5): 517-32.

Hayashi, K., Momoi, Y., Tanuma, N., Kishimoto, A., Ogoh, H., Kato, H., Suzuki, M., Sakamoto, Y., Inoue, Y., Nomura, M., Kiyonari, H., Sakayori, M., Fukamachi, K., Kakugawa, Y., Yamashita, Y., Ito, S., Sato, I., Suzuki, A., Nishio, M., Suganuma, M., Watanabe, T. and Shima, H. (2015) Abrogation of protein phosphatase 6 promotes skin carcinogenesis induced by DMBA. *Oncogene* 34(35): 4647-55.

Iezaki, T., Onishi, Y., Ozaki, K., Fukasawa, K., Takahata, Y., Nakamura, Y., Fujikawa, K., Takarada, T., Yoneda, Y., Yamashita, Y., Shioi, G. and Hinoi, E. (2015) The Transcriptional Modulator Interferon-Related Developmental Regulator 1 in Osteoblasts Suppresses Bone Formation and Promotes Bone Resorption. *J Bone Miner Res.* 2015 Sep 22. doi: 10.1002/jbmr.2720.

Dong, B., Endo, I., Ohnishi, Y., Kondo, T., Hasegawa, T., Amizuka, N., Kiyonari, H., Shioi, G., Abe, M., Fukumoto, S. and Matsumoto, T. (2015) Calcilytic Ameliorates Abnormalities of Mutant Calcium-Sensing Receptor (CaSR) Knock-In Mice Mimicking Autosomal Dominant Hypocalcemia (ADH). *J Bone Miner Res.* 2015 May 12. doi: 10.1002/jbmr.2551.

Hashidate-Yoshida, T., Harayama, T., Hishikawa, D., Morimoto, R., Hamano, F., Tokuoka, S. M., Eto, M., Tamura-Nakano, M., Yanobu-Takanashi, R., Mukumoto, Y., Kiyonari, H., Okamura, T., Kita, Y., Shindou, H. and Shimizu, T. (2015) Fatty acid remodeling by LPCAT3 enriches arachidonate in phospholipid membranes and regulates triglyceride transport. *Elife* 4. doi: 10.7554/eLife.06328.

Mukai, S., Yabuta, N., Yoshida, K., Okamoto, A., Miura, D., Furuta, Y., Abe, T. and Nojima H. (2015) Lats1 suppresses centrosome overduplication by modulating the stability of Cdc25B. *Sci Rep* 5:16173. doi: 10.1038/srep16173.

Irie, M., Yoshikawa, M., Ono, R., Iwafune, H., Furuse, T., Yamada, I., Wakana, S., Yamashita, Y., Abe, T., Ishino, F., Kaneko-Ishino, T. (2015) Cognitive Function Related to the Sirh11/Zcchc16 Gene Acquired from an LTR Retrotransposon in Eutherians, *PLoS Genet* 11(9):e1005521.

Arimoto K, Hishiki T, Kiyonari H, Abe T, Cheng C, Yan M, Fan JB, Futakuchi M, Tsuda H, Murakami Y, Suzuki H, Zhang DE, Shimotohno K. (2015) Murine Herc6 Plays a Critical Role in Protein ISGylation In Vivo and Has an ISGylation-Independent Function in Seminal Vesicles. *J Interferon Cytokine Res.* 35(5):351-8.

## 2014

Oki, T., Nishimura, K., Kitaura, J., Togami, K., Maehara, A., Izawa, K., Sakaue-Sawano, A., Niida, A., Miyano, S., Aburatani, H., Kiyonari, H., Miyawaki, A. and Kitamura, T. (2014) A novel cell-cycle-indicator, mVenus-p27K-, identifies quiescent cells and visualizes G0-G1 transition. *Sci Rep* 4: 4012.

Kawaue, T., Sagou, K., Kiyonari, H., Ota, K., Okamoto, M., Shinoda, T., Kawaguchi, A. and Miyata, T. (2014) Neurogenin2-d4Venus and Gadd45g-d4Venus transgenic mice: visualizing mitotic and migratory behaviors of cells committed to the neuronal lineage in the developing mammalian brain. *Dev Growth Differ* 56(4): 293-304.

Watanabe, S., De Zan, T., Ishizaki, T., Yasuda, S., Kamijo, H., Yamada, D., Aoki, T., Kiyonari, H., Kaneko, H., Shimizu, R., Yamamoto, M., Goshima, G. and Narumiya, S. (2013) Loss of a Rho-regulated actin nucleator, mDia2, impairs cytokinesis during mouse fetal erythropoiesis. *Cell Rep* 5(4): 926-32.

Goriki, A., Hatanaka, F., Myung, J., Kim, J. K., Yoritaka, T., Tanoue, S., Abe, T., Kiyonari, H., Fujimoto, K., Kato, Y., Todo, T., Matsubara, A., Forger, D. and Takumi, T. (2014) A novel protein, CHRONO, functions as a core component of the mammalian circadian clock. *PLoS Biol* 12(4): e1001839.

Hayashi, S., Inoue, Y., Kiyonari, H., Abe, T., Misaki, K., Moriguchi, H., Tanaka, Y. and Takeichi, M. (2014) Protocadherin-17 mediates collective axon extension by recruiting actin regulator complexes to interaxonal contacts. *Dev Cell* 30(6): 673-87.

Inoue, T., Hoshina, N., Nakazawa, T., Kiyama, Y., Kobayashi, S., Abe, T., Yamamoto, T., Manabe, T. and Yamamoto, T. (2014) LMTK3 deficiency causes pronounced locomotor hyperactivity and impairs endocytic trafficking. *J Neurosci* 34(17): 5927-37.

Kitamura, A., Sasaki, Y., Abe, T., Kano, H. and Yasutomo, K. (2014) An inherited mutation in NLRC4 causes autoinflammation in human and mice. *J Exp Med* 211(12): 2385-96.



Miyazaki, H., Oyama, F., Inoue, R., Aosaki, T., Abe, T., Kiyonari, H., Kino, Y., Kurosawa, M., Shimizu, J., Ogiwara, I., Yamakawa, K., Koshimizu, Y., Fujiyama, F., Kaneko, T., Shimizu, H., Nagatomo, K., Yamada, K., Shimogori, T., Hattori, N., Miura, M. and Nukina, N. (2014) Singular localization of sodium channel beta4 subunit in unmyelinated fibres and its role in the striatum. *Nat Commun* 5: 5525.

Nakahata, S., Ichikawa, T., Maneesaay, P., Saito, Y., Nagai, K., Tamura, T., Manachai, N., Yamakawa, N., Hamasaki, M., Kitabayashi, I. Arai, Y., Kanai, Y., Taki, T., Abe, T., Kiyonari, H., Shimoda, K., Ohshima, K., Horii, A., Shima, H., Taniwaki, M., Yamaguchi, R. and Morishita, K. (2014) Loss of NDRG2 expression activates PI3K-AKT signalling via PTEN phosphorylation in ATLL and other cancers. *Nat Commun* 5: 3393.

Ohkubo, N., Matsubara, E., Yamanouchi, J., Akazawa, R., Aoto, M., Suzuki, Y., Sakai, I., Abe, T., Kiyonari, H., Matsuda, S., Yasukawa, M. and Mitsuda, N. (2014) Abnormal behaviors and developmental disorder of hippocampus in zinc finger protein 521 (ZFP521) mutant mice. *PLoS One* 9(3): e92848.

Okumura, S., Fujita, T., Cai, W., Jin, M., Namekata, I., Mototani, Y., Jin, H., Ohnuki, Y., Tsuneoka, Y., Kurotani, R. Suita, K., Kawakami, Y., Hamaguchi, S., Abe, T., Kiyonari, H., Tsunematsu, T., Bai, Y., Suzuki, S., Hidaka, Y., Umemura, M., Ichikawa, Y., Yokoyama, U., Sato, M., Ishikawa, F., Izumi-Nakaseko, H., Adachi-Akahane, S., Tanaka, H. and Ishikawa, Y. (2014) Epac1-dependent phospholamban phosphorylation mediates the cardiac response to stresses. *J Clin Invest* 124(6): 2785-801.

Susaki, E. A., Tainaka, K., Perrin, D., Kishino, F., Tawara, T., Watanabe, T. M., Yokoyama, C., Onoe, H., Eguchi, M., Yamaguchi, S., Abe, T., Kiyonari, H., Shimizu, Y., Miyawaki, A., Yokota, H. and Ueda, HR. (2014) Whole-brain imaging with single-cell resolution using chemical cocktails and computational analysis. *Cell* 157(3): 726-39.

Tamai, S., Imaizumi, K., Kurabayashi, N., Nguyen, M. D., Abe, T., Inoue, M., Fukada,

Y. and Sanada, K. (2014) Neuroprotective role of the basic leucine zipper transcription factor NFIL3 in models of amyotrophic lateral sclerosis. *J Biol Chem* 289(3): 1629-38.

Yano, K., Carter, C., Yoshida, N., Abe, T., Yamada, A., Nitta, T., Ishimaru, N., Takada, K., Butcher, G. W. and Takahama, Y. (2014) Gimap3 and Gimap5 cooperate to maintain T-cell numbers in the mouse. *Eur J Immunol* 44(2): 561-72.

Morimoto, A., Tomita, S., Imanishi, M., Shioi, G., Kihira, Y., Izawa-Ishizawa, Y., Takaku, M., Hashimoto, I., Ikeda, Y., Nakanishi, H. and Tamaki, T. (2014) Overexpressed HIF-2alpha in Endothelial Cells Promotes Vascularization and Improves Random Pattern Skin Flap Survival. *Plast Reconstr Surg Glob Open* 2(4): e132.

Ishiguro, K., Kim, J., Shibuya, H., Hernandez-Hernandez, A., Suzuki, A., Fukagawa, T., Shioi, G., Kiyonari, H., Li, X. C., Schimenti, J., Höög, C. and Watanabe, Y. (2014) Meiosis-specific cohesin mediates homolog recognition in mouse spermatocytes. *Genes Dev* 28(6): 594-607.

Hirano, A., Kurabayashi, N., Nakagawa, T., Shioi, G., Todo, T., Hirota, T. and Fukada, Y. (2014) In vivo role of phosphorylation of cryptochrome 2 in the mouse circadian clock. *Mol Cell Biol* 34(24): 4464-73.

Okumura S, Fujita T, Cai W, Jin M, Namekata I, Mototani Y, Jin H, Ohnuki Y, Tsuneoka Y, Kurotani R, Suita K, Kawakami Y, Hamaguchi S, Abe T, Kiyonari H, Tsunematsu T, Bai Y, Suzuki S, Hidaka Y, Umemura M, Ichikawa Y, Yokoyama U, Sato M, Ishikawa F, Izumi-Nakaseko H, Adachi-Akahane S, Tanaka H, Ishikawa Y. (2014) Epac1-dependent hospholamban phosphorylation mediates the cardiac response to stresses. *J Clin Invest*. 124(6):2785-2801. doi: 10.1172/JCI64784.

## 2013

Abe, T. and Fujimori, T. (2013) Reporter mouse lines for fluorescent imaging. *Dev Growth Differ* 55(4): 390-405.

Abe, T., Aizawa, S. and Fujimori, T. (2013) Live imaging of early mouse embryos using fluorescently labeled transgenic mice. *Methods Mol Biol* 1052: 101-8. doi: 10.1007/7651\_2013\_20.

Abe, T., Sakaue-Sawano, A., Kiyonari, H., Shioi, G., Inoue, K., Horiuchi, T., Nakao, K., Miyawaki, A., Aizawa, S. and Fujimori, T. (2013) Visualization of cell cycle in mouse embryos with Fucci2 reporter directed by Rosa26 promoter. *Development* 140(1): 237-46.

Fujikura, K., Setsu, T., Tanigaki, K., Abe, T., Kiyonari, H., Terashima, T. and Sakisaka, T. (2013) Kif14 mutation causes severe brain malformation and hypomyelination. *PLoS One* 8(1): e53490.

Ikeshima-Kataoka, H., Abe, Y., Abe, T. and Yasui, M. (2013) Immunological function of aquaporin-4 in stab-wounded mouse brain in concert with a pro-inflammatory cytokine inducer, osteopontin. *Mol Cell Neurosci*. 56:65-75. doi: 10.1016/j.mcn.2013.02.002.

Tanaka, S., Kawaguchi, S. Y., Shioi, G. and Hirano, T. (2013) Long-term potentiation of inhibitory synaptic transmission onto cerebellar Purkinje neurons contributes to adaptation of vestibulo-ocular reflex. *J Neurosci* 33(43): 17209-20.

Park, S., Shcheynikov, N., Hong, J. H., Zheng, C., Suh, S. H., Kawaai, K., Ando, H., Mizutani, A., Abe, T., Kiyonari, H., Seki, G., Yule, D., Mikoshiba, K. and Muallem, S. (2013) Irbit mediates synergy between ca(2+) and cAMP signaling pathways during epithelial transport in mice. *Gastroenterology* 145(1): 232-41.

Sakaguchi, M., Sharmin, S., Taguchi, A., Ohmori, T., Fujimura, S., Abe, T., Kiyonari,

H., Komatsu, Y., Mishina, Y., Asashima, M., Araki, E. and Nishinakamura, R. (2013) The phosphatase Dullard negatively regulates BMP signalling and is essential for nephron maintenance after birth. *Nat Commun* 4: 1398.

Takao, D., Nemoto, T., Abe, T., Kiyonari, H., Kajiura-Kobayashi, H., Shiratori, H. and Nonaka, S. (2013) Asymmetric distribution of dynamic calcium signals in the node of mouse embryo during left-right axis formation. *Dev Biol* 376(1): 23-30.

Takarada, T., Hinoi, E., Nakazato, R., Ochi, H., Xu, C., Tsuchikane, A., Takeda, S., Karsenty, G., Abe, T., Kiyonari, H. and Yoneda, Y. (2013) An analysis of skeletal development in osteoblast-specific and chondrocyte-specific runt-related transcription factor-2 (Runx2) knockout mice. *J Bone Miner Res* 28(10): 2064-9.

Tamaki, M., Fujitani, Y., Hara, A., Uchida, T., Tamura, Y., Takeno, K., Kawaguchi, M., Watanabe, T., Ogihara, T., Fukunaka, A., Shimizu, T., Mita, T., Kanazawa, A., Imaizumi, M. O., Abe, T., Kiyonari, H., Hojyo, S., Fukada, T., Kawauchi, T., Nagamatsu, S., Hirano, T., Kawamori, R. and Watada, H. (2013) The diabetes-susceptible gene SLC30A8/ZnT8 regulates hepatic insulin clearance. *J Clin Invest* 123(10): 4513-24.

Tanaka, S. S., Nakane, A., Yamaguchi, Y. L., Terabayashi, T., Abe, T., Nakao, K., Asashima, M., Steiner, K. A., Tam, P. P. and Nishinakamura, R. (2013) Dullard/Ctdnep1 Modulates WNT Signalling Activity for the Formation of Primordial Germ Cells in the Mouse Embryo. *PLoS One* 8(3): e57428.

Yagi, H., Nakagawa, N., Saito, T., Kiyonari, H., Abe, T., Toda, T., Wu, S. W., Khoo, K. H., Oka, S. and Kato, K. (2013) AGO61-dependent GlcNAc modification primes the formation of functional glycans on alpha-dystroglycan. *Sci Rep* 3: 3288.

Nakashima, A., Takeuchi, H., Imai, T., Saito, H., Kiyonari, H., Abe, T., Chen, M., Weinstein, L. S., Yu, C. R., Storm, D. R., Nishizumi, H. and Sakano, H. (2013) Agonist-independent GPCR activity regulates anterior-posterior targeting of olfactory sensory neurons. *Cell* 154(6): 1314-25.

Masuda, K., Ripley, B., Nishimura, R., Mino, T., Takeuchi, O., Shioi, G., Kiyonari, H. and Kishimoto, T. (2013) Arid5a controls IL-6 mRNA stability, which contributes to elevation of IL-6 level in vivo. *Proc Natl Acad Sci U S A* 110(23): 9409-14.

Lee, J., Kim, K. E., Choi, D. K., Jang, J. Y., Jung, J. J., Kiyonari, H., Shioi, G., Chang, W., Suda, T., Mochizuki, N., Nakaoka, Y., Komuro, I., Yoo, O.J. and Koh, G.Y. (2013) Angiopoietin-1 guides directional angiogenesis through integrin  $\alpha$ v $\beta$ 5 signaling for recovery of ischemic retinopathy. *Sci Transl Med* 5(203): 203ra127.

Kimura, I., Ozawa, K., Inoue, D., Imamura, T., Kimura, K., Maeda, T., Terasawa, K., Kashihara, D., Hirano, K., Tani, T., Takahashi, T., Miyauchi, S., Shioi, G., Inoue, H. and Tsujimoto, G. (2013) The gut microbiota suppresses insulin-mediated fat accumulation via the short-chain fatty acid receptor GPR43. *Nat Commun* 4: 1829. doi: 10.1038/ncomms2852.

Huong le, T., Kobayashi, M., Nakata, M., Shioi, G., Miyachi, H., Honjo, T. and Nagaoka, H. (2013) In vivo analysis of Aicda gene regulation: a critical balance between upstream enhancers and intronic silencers governs appropriate expression. *PLoS One* 8(4): e61433.

Matsuo, Y., Irie, K., Kiyonari, H., Okuyama, H., Nakamura, H., Son, A., Lopez-Ramos, D. A., Tian, H., Oka, S., Okawa, K., Kizaka-Kondoh, S., Masutani, H., Yodoi, J. (2013) The protective role of the transmembrane thioredoxin-related protein TMX in inflammatory liver injury. *Antioxid Redox Signal* 18(11): 1263-72.

Kon, S., Minegishi, N., Tanabe, K., Watanabe, T., Funaki, T., Wong, W. F., Sakamoto, D., Higuchi, Y., Kiyonari, H., Asano, K., Iwakura, Y., Fukumoto, M., Osato, M., Sanada, M., Ogawa, S., Nakamura, T. and Satake, M. (2013) Smap1 deficiency perturbs receptor trafficking and predisposes mice to myelodysplasia. *J Clin Invest* 123(3): 1123-37.

Imuta, Y., Kiyonari, H., Jang, C. W., Behringer, R. R. and Sasaki, H. (2013) Generation

of knock-in mice that express nuclear enhanced green fluorescent protein and tamoxifen-inducible Cre recombinase in the notochord from *Foxa2* and *T* loci. *Genesis* 51(3): 210-8.

Hoshina, N., Tanimura, A., Yamasaki, M., Inoue, T., Fukabori, R., Kuroda, T., Yokoyama, K., Tezuka, T., Sagara, H., Hirano, S., Kiyonari, H., Takada, M., Kobayashi, K., Watanabe, M., Kano, M., Nakazawa, T., Yamamoto, T. (2013) Protocadherin 17 regulates presynaptic assembly in topographic corticobasal Ganglia circuits. *Neuron* 78(5): 839-54.

Tsujino, K., Narumi, R., Masumoto, KH., Susaki, EA., Shinohara, Y., Abe, T., Iigo, M., Wada, A., Nagano, M., Shigeyoshi, Y., Ueda, HR. (2013) Establishment of TSH  $\beta$  real-time monitoring system in mammalian photoperiodism. *Genes Cells* 18(7): 575-588.

Matsuo, Y., Irie, K., Kiyonari, H., Okuyama, H., Nakamura, H., Son, A., Lopez-Ramos, DA., Tian, H., Oka, S., Okawa, K., Kizaka-Kondoh, S., Masutani, H. and Yodoi, J. (2013) The protective role of transmembrane thioredoxin-related protein TMX in inflammatory liver injury. *Antioxid Redox Signal* 18(11): 1263-72.

## 2012

Ohmura, T., Shioi, G., Hirano, M. and Aizawa, S. (2012) Neural tube defects by NUA1 and NUA2 double mutation. *Dev Dyn* 241(8): 1350-64.

Tao, H., Inoue, K., Kiyonari, H., Bassuk, A. G., Axelrod, J. D., Sasaki, H., Aizawa, S. and Ueno, N. (2012) Nuclear localization of Prickle2 is required to establish cell polarity during early mouse embryogenesis. *Dev Biol* 364(2): 138-48.

Shinohara, R., Thumkeo, D., Kamijo, H., Kaneko, N., Sawamoto, K., Watanabe, K., Takebayashi, H., Kiyonari, H., Ishizaki, T., Furuyashiki, T. and Narumiya, S. (2012) A role for mDia, a Rho-regulated actin nucleator, in tangential migration of interneuron precursors. *Nat Neurosci* 15(3): 373-80.

Okuda-Ashitaka, E., Minami, T., Tsubouchi, S., Kiyonari, H., Iwamatsu, A., Noda, T., Handa, H. and Ito, S. (2012) Identification of NIPSNAP1 as a nocistatin-interacting protein involving pain transmission. *J Biol Chem* 287(13): 10403-13.

Kogo, H., Tsutsumi, M., Inagaki, H., Ohye, T., Kiyonari, H. and Kurahashi, H. (2012) HORMAD2 is essential for synapsis surveillance during meiotic prophase via the recruitment of ATR activity. *Genes Cells* 17(11): 897-912.

Iwano, T., Masuda, A., Kiyonari, H., Enomoto, H. and Matsuzaki, F. (2012) 'Prox1 postmitotically defines dentate gyrus cells by specifying granule cell identity over CA3 pyramidal cell fate in the hippocampus', *Development* 139(16): 3051-62.

Sato, S., Ikeda, K., Shioi, G., Nakao, K., Yajima, H. and Kawakami, K. (2012) Regulation of Six1 expression by evolutionarily conserved enhancers in tetrapods. *Dev Biol* 368(1): 95-108.

Nishimura, R., Wakabayashi, M., Hata, K., Matsubara, T., Honma, S., Wakisaka, S., Kiyonari, H., Shioi, G., Yamaguchi, A., Tsumaki, N., Akiyama, H. and Yoneda, T. (2012) Osterix regulates calcification and degradation of chondrogenic matrices through

matrix metalloproteinase 13 (MMP13) expression in association with transcription factor Runx2 during endochondral ossification. *J Biol Chem* 287(40): 33179-90.

Nakagawa, S., Ip, J. Y., Shioi, G., Tripathi, V., Zong, X., Hirose, T. and Prasanth, K. V. (2012) Malat1 is not an essential component of nuclear speckles in mice. *RNA* 18(8): 1487-99.

Kanemaru, K., Nakamura, Y., Sato, K., Kojima, R., Takahashi, S., Yamaguchi, M., Ichinohe, M., Kiyonari, H., Shioi, G., Kabashima, K., Nakahigashi, K., Asagiri, M., Jamora, C., Yamaguchi, H. and Fukami, K. (2012) Epidermal phospholipase C $\delta$ 1 regulates granulocyte counts and systemic interleukin-17 levels in mice. *Nat Commun* 3: 963. doi: 10.1038/ncomms1960.

Inazuka, F., Sugiyama, N., Tomita, M., Abe, T., Shioi, G. and Esumi, H. (2012) Muscle-specific knock-out of NUA family SNF1-like kinase 1 (NUAK1) prevents high fat diet-induced glucose intolerance. *J Biol Chem* 287(20): 16379-89.

Aoyama, K., Wang, F., Matsumura, N., Kiyonari, H., Shioi, G., Tanaka, K., Kinoshita, C., Kikuchi-Utsumi, K., Watabe, M. and Nakaki, T. (2012) Increased neuronal glutathione and neuroprotection in GTRAP3-18-deficient mice. *Neurobiol Dis* 45(3): 973-82.

Izawa, K., Yamanishi, Y., Maehara, A., Takahashi, M., Isobe, M., Ito, S., Kaitani, A., Matsukawa, T., Matsuoka, T., Nakahara, F., Oki, T., Kiyonari, H., Abe, T., Okumura, K., Kitamura, T. and Kitaura, J. (2012) The Receptor LMIR3 Negatively Regulates Mast Cell Activation and Allergic Responses by Binding to Extracellular Ceramide. *Immunity* 37(5): 827-39.

Niisato, E., Nagai, J., Yamashita, N., Abe, T., Kiyonari, H., Goshima, Y. and Ohshima, T. (2012) CRMP4 suppresses apical dendrite bifurcation of CA1 pyramidal neurons in the mouse hippocampus. *Dev Neurobiol* 72(11): 1447-57.

Sugimoto, M., Kondo, M., Hirose, M., Suzuki, M., Mekada, K., Abe, T., Kiyonari, H.,



Ogura, A., Takagi, N., Artzt, K. and Abe, K. (2012) Molecular Identification of t(w5): Vps52 Promotes Pluripotential Cell Differentiation through Cell-Cell Interactions. *Cell Rep* 2(5): 1363-74.

Fukuhara, S., Simmons, S., Kawamura, S., Inoue, A., Orba, Y., Tokudome, T., Sunden, Y., Arai, Y., Moriwaki, K., Ishida, J., uemura, A., Kiyonari, H., Abe, T., Fukamizu, A., Hirashima, M., Sawa, H., Aoki, J., Ishii, M. and Mochizuki, N. (2012) The sphingosine-1-phosphate transporter Spns2 expressed on endothelial cells regulates lymphocyte trafficking in mice. *J Clin Invest* 122(4): 1416-26.

Nakae, J., Cao, Y., Hakuno, F., Takemori, H., Kawano, Y., Sekioka, R., Abe, T., Kiyonari, H., Tanaka, T., Sakai, J., Takahashi, S. and Itoh, H. (2012) Novel repressor regulates insulin sensitivity through interaction with Foxo1. *EMBO J* 31(10): 2275-95.

Kogo, H., Tsutsumi, M., Ohye, T., Inagaki, H., Abe, T. and Kurahashi, H. (2012) HORMAD1-dependent checkpoint/surveillance mechanism eliminates asynaptic oocytes. *Genes Cells* 17(6): 439-54.

Konno, D., Iwashita, M., Satoh, Y., Momiyama, A., Abe, T., Kiyonari, H. and Matsuzaki, F. (2012) The Mammalian DM Domain Transcription Factor Dmrta2 Is Required for Early Embryonic Development of the Cerebral Cortex. *PLoS One* 7(10): e46577.

Suzuki, M., Tanaka, H., Tanimura, A., Tanabe, K., Oe, N., Rai, S., Kon, S., Fukumoto, M., Takei, K., Abe, T., Matsumura, I., Kanakura, Y. and Watanabe, T. (2012) The clathrin assembly protein PICALM is required for erythroid maturation and transferrin internalization in mice. *PLoS One* 7(2): e31854.

## 2011

Shioi, G., Kiyonari, H., Abe, T., Nakao, K., Fujimori, T., Jang, C. W., Huang, C. C., Akiyama, H., Behringer, R. R. and Aizawa, S. (2011) A mouse reporter line to conditionally mark nuclei and cell membranes for in vivo live-imaging. *Genesis* 49(7): 570-8.

Abe, T., Kiyonari, H., Shioi, G., Inoue, K., Nakao, K., Aizawa, S. and Fujimori, T. (2011) Establishment of conditional reporter mouse lines at ROSA26 locus for live cell imaging. *Genesis* 49(7): 579-90.

Nakagawa, S., Naganuma, T., Shioi, G. and Hirose, T. (2011) Paraspeckles are subpopulation-specific nuclear bodies that are not essential in mice. *J Cell Biol* 193(1): 31-9.

Hara, T., Mimura, K., Abe, T., Shioi, G., Seiki, M. and Sakamoto, T. (2011) Deletion of the *Mint3/Apba3* gene in mice abrogates macrophage functions and increases resistance to lipopolysaccharide-induced septic shock. *J Biol Chem* 286(37): 32542-51.

Ishikawa, T., Gotoh, N., Murayama, C., Abe, T., Iwashita, M., Matsuzaki, F., Suzuki, T. and Yamamoto, T. (2011) IgSF molecule MDGA1 is involved in radial migration and positioning of a subset of cortical upper-layer neurons. *Dev Dyn* 240(1): 96-107.

Kobayashi, M., Sabouri, Z., Sabouri, S., Kitawaki, Y., Pommier, Y., Abe, T., Kiyonari, H. and Honjo, T. (2011) Decrease in topoisomerase I is responsible for activation-induced cytidine deaminase (AID)-dependent somatic hypermutation. *Proc Natl Acad Sci U S A* 108(48): 19305-10.

Kuroda, K., Yamada, S., Tanaka, M., Iizuka, M., Yano, H., Mori, D., Tsuboi, D., Nishioka, T., Namba, T., Iizuka, Y., Kubota, S., Nagai, T., Ibi, D., Wang, R., Enomoto, A., Isotani-Sakakibara, M., Asai, N., Kimura, K., Kiyonari, H., Abe, T., Mizoguchi, A., Sokabe, M., Takahashi, M., Yamada, K. and Kaibuchi, K. (2011) Behavioral alterations associated with targeted disruption of exons 2 and 3 of the *Disc1* gene in the mouse.

Hum Mol Genet 20(23): 4666-83.

Mouri, Y., Yano, M., Shinzawa, M., Shimo, Y., Hirota, F., Nishikawa, Y., Nii, T., Kiyonari, H., Abe, T., Uehara, H., Izumi, K., Tamada, K., Chen, L., Penninger, J.M., Inoue, J., Akiyama, T. and Matsumoto, M. (2011) Lymphotoxin signal promotes thymic organogenesis by eliciting RANK expression in the embryonic thymic stroma. *J Immunol* 186(9): 5047-57.

Sano, T., Kim, Y. J., Oshima, E., Shimizu, C., Kiyonari, H., Abe, T., Higashi, H., Yamada, K. and Hirabayashi, Y. (2011) Comparative characterization of GPRC5B and GPRC5CLacZ knockin mice; behavioral abnormalities in GPRC5B-deficient mice. *Biochem Biophys Res Commun* 412(3): 460-5.

Sato, Y., Iketani, M., Kurihara, Y., Yamaguchi, M., Yamashita, N., Nakamura, F., Arie, Y., Kawasaki, T., Hirata, T., Abe, T., Kiyonari, H., Strittmatter, S.M., Goshima, Y. and Takei, K. (2011) Cartilage acidic protein-1B (LOTUS), an endogenous Nogo receptor antagonist for axon tract formation. *Science* 333(6043): 769-73.

Shibata, M., Nakao, H., Kiyonari, H., Abe, T. and Aizawa, S. (2011) MicroRNA-9 regulates neurogenesis in mouse telencephalon by targeting multiple transcription factors. *J Neurosci* 31(9): 3407-22.

Takahashi, M., Okimura, Y., Iguchi, G., Nishizawa, H., Yamamoto, M., Suda, K., Kitazawa, R., Fujimoto, W., Takahashi, K., Zolotaryov, F. N., Hong, K. S., Kiyonari, H., Abe, T., Kaji, H., Kitazawa, S., Kasuga, M., Chihara, K. and Takahashi, Y. (2011) Chemerin regulates beta-cell function in mice. *Sci Rep* 1: 123. doi: 10.1038/srep00123.

Tao, H., Manak, J. R., Sowers, L., Mei, X., Kiyonari, H., Abe, T., Dahdaleh, N. S., Yang, T., Wu, S., Chen, S., Fox, M.H., Gurnett, C., Montine, T., Bird, T., Shaffer, L.G., Rosenfeld, J.A., McConnell, J., Madan-Khetarpal, S., Berry-Kravis, E., Griesbach, H., Saneto, R.P., Scott, M.P., Antic, D., Reed, J., Boland, R., Ehaideb, S.N., El-Shanti, H., Mahajan, V.B., Ferguson, P.J., Axelrod, J.D., Lehesjoki, A.E., Fritsch, B., Slusarski, D.C., Wemmie, J., Ueno, N. and Bassuk, A.G. (2011) Mutations in prickle orthologs

cause seizures in flies, mice, and humans. *Am J Hum Genet* 88(2): 138-49.

Yabuta, Y., Ohta, H., Abe, T., Kurimoto, K., Chuma, S. and Saitou, M. (2011) TDRD5 is required for retrotransposon silencing, chromatoid body assembly, and spermiogenesis in mice. *J Cell Biol* 192(5): 781-95.

Yamanaka, Y., Kitano, A., Takao, K., Prasansuklab, A., Mushiroda, T., Yamazaki, K., Kumada, T., Shibata, M., Takaoka, Y., Awaya, T., Kato, T., Abe, T., Iwata, N., Miyakawa, T., Nakamura, Y., Nakahata, T. and Heike, T. (2011) Inactivation of fibroblast growth factor binding protein 3 causes anxiety-related behaviors. *Mol Cell Neurosci* 46(1): 200-12.

Zhang, J., Eto, K., Honmyou, A., Nakao, K., Kiyonari, H. and Abe, S. (2011) Neuregulins are essential for spermatogonial proliferation and meiotic initiation in neonatal mouse testis. *Development* 138(15): 3159-68.

Yamaguchi, Y. L., Tanaka, S. S., Oshima, N., Kiyonari, H., Asashima, M. and Nishinakamura, R. (2011) Translocon-associated protein subunit Trap-gamma/Ssr3 is required for vascular network formation in the mouse placenta. *Dev Dyn* 240(2): 394-403.

Kamiya, D., Banno, S., Sasai, N., Ohgushi, M., Inomata, H., Watanabe, K., Kawada, M., Yakura, R., Kiyonari, H., Nakao, K., Jakt, L.M., Nishikawa, S. and Sasai, Y. (2011) Intrinsic transition of embryonic stem-cell differentiation into neural progenitors. *Nature* 470(7335): 503-9.

Inoue, Y., Tashiro-Yamaji, J., Hayashi, M., Kiyonari, H., Shimizu, T., Ibata, M., Yamana, H., Kubota, T., Tanigawa, N. and Yoshida, R. (2011) Transgene number-dependent, gene expression rate-independent rejection of Dd -, Kd-, or Dd Kd -transgenic mouse skin or tumor cells from C57BL/6 Db Kb mice. *Microbiol Immunol* 55(6): 446-53.

## 2010

Nakao, K. and Inoue, K. (2010) Cryopreservation for Broader Production of Transgenic Mice by DNA Injection into Zygotes. *Exp Anim* 59(2): 225-9.

Kiyonari, H., Kaneko, M., Abe, S. and Aizawa, S. (2010) Three inhibitors of FGF receptor, ERK, and GSK3 establishes germline-competent embryonic stem cells of C57BL/6N mouse strain with high efficiency and stability. *Genesis* 48(5): 317-27.

Uchiyama, Y., Sakaguchi, M., Terabayashi, T., Inenaga, T., Inoue, S., Kobayashi, C., Oshima, N., Kiyonari, H., Nakagata, N., Sato, Y., Sekiguchi, K., Miki, H., Araki, E., Fujimura, S., Tanaka, SS. and Nishinakamura, R. (2010) Kif26b, a kinesin family gene, regulates adhesion of the embryonic kidney mesenchyme. *Proc Natl Acad Sci U S A* 107(20): 9240-5.

Sakurai, Y., Kurokawa, D., Kiyonari, H., Kajikawa, E., Suda, Y. and Aizawa, S. (2010) Otx2 and Otx1 protect diencephalon and mesencephalon from caudalization into metencephalon during early brain regionalization. *Dev Biol* 347(2): 392-403.

Kanemaru, K., Nakahara, M., Nakamura, Y., Hashiguchi, Y., Kouchi, Z., Yamaguchi, H., Oshima, N., Kiyonari, H. and Fukami, K. (2010) Phospholipase C- $\epsilon$ 2 is highly expressed in the habenula and retina. *Gene Expr Patterns* 10(2-3): 119-26.

Chandana, E. P., Maeda, Y., Ueda, A., Kiyonari, H., Oshima, N., Yamamoto, M., Kondo, S., Oh, J., Takahashi, R., Yoshida, Y., Kawashima, S., Alexander, D.B., Kitayama, H., Takahashi, C., Tabata, Y., Matsuzaki, T. and Noda, M. (2010) Involvement of the Reck tumor suppressor protein in maternal and embryonic vascular remodeling in mice. *BMC Dev Biol* 10: 84. doi: 10.1186/1471-213X-10-84.

Sato, S., Ikeda, K., Shioi, G., Ochi, H., Ogino, H., Yajima, H. and Kawakami, K. (2010) Conserved expression of mouse Six1 in the pre-placodal region (PPR) and identification of an enhancer for the rostral PPR. *Dev Biol* 344(1): 158-71.

Ito, A., Shinmyo, Y., Abe, T., Oshima, N., Tanaka, H. and Ohta, K. (2010) Tsukushi is required for anterior commissure formation in mouse brain. *Biochem Biophys Res Commun* 402(4): 813-8.

Kumada, T., Yamanaka, Y., Kitano, A., Shibata, M., Awaya, T., Kato, T., Okawa, K., Abe, T., Oshima, N., Nakahata, T. and Heike, T. (2010) Ttyh1, a Ca(2+)-binding protein localized to the endoplasmic reticulum, is required for early embryonic development. *Dev Dyn* 239(8): 2233-45.

Oboki, K., Ohno, T., Kajiwara, N., Arae, K., Morita, H., Ishii, A., Nambu, A., Abe, T., Kiyonari, H., Matsumoto, K., Sudo, K., Okumura, K., Saito, H. and Nakae, S. (2010) IL-33 is a crucial amplifier of innate rather than acquired immunity. *Proc Natl Acad Sci U S A* 107(43): 18581-6.

Tanabe, K., Kani, S., Shimizu, T., Bae, Y. K., Abe, T. and Hibi, M. (2010) Atypical protein kinase C regulates primary dendrite specification of cerebellar Purkinje cells by localizing Golgi apparatus. *J Neurosci* 30(50): 16983-92.

Yamanishi, Y., Kitaura, J., Izawa, K., Kaitani, A., Komeno, Y., Nakamura, M., Yamazaki, S., Enomoto, Y., Oki, T., Akiba, H., Abe, T., Komori, T., Morikawa, Y., Kiyonari, H., Takai, T., Okumura, K. and Kitamura, T. (2010) TIM1 is an endogenous ligand for LMIR5/CD300b: LMIR5 deficiency ameliorates mouse kidney ischemia/reperfusion injury. *J Exp Med* 207(7): 1501-11.

Suzuki, S., Yokoyama, U., Abe, T., Kiyonari, H., Yamashita, N., Kato, Y., Kurotani, R., Sato, M., Okumura, S. and Ishikawa, Y. (2010) Differential roles of Epac in regulating cell death in neuronal and myocardial cells. *J Biol Chem* 285(31): 24248-59.

## 2009

Yamakoshi, K., Takahashi, A., Hirota, F., Nakayama, R., Ishimaru, N., Kubo, Y., Mann, D.J., Ohmura, M., Hirao, A., Saya, H., Arase, S., Hayashi, Y., Nakao, K., Matsumoto, M., Ohtani, N. and Hara, E. (2009) Real-time in vivo imaging of p16Ink4a reveals cross-talk with p53. *J Cell Biol* 186(3): 393-407.

Ukita, K., Hirahara, S., Oshima, N., Imuta, Y., Yoshimoto, A., Jang, C.W., Oginuma, M., Saga, Y., Behringer, R.R., Kondoh, H. and Sasaki, H. (2009) Wnt signaling maintains the notochord fate for progenitor cells and supports the posterior extension of the notochord. *Mech Dev* 126(10): 791-803.

Okuyama, H., Yoshida, T., Son, A., Oka, S., Wang, D., Nakayama, R., Masutani, H., Nakamura, H., Nabeshima, Y. and Yodoi, J. (2009) Thioredoxin binding protein 2 modulates natural killer T cell-dependent innate immunity in the liver: possible link to lipid metabolism. *Antioxid Redox Signal*. 11(10): 2585-93.

Kato, M., Hasunuma, N., Manabe, M., Nakayama, R., Takeda, J., Itami, S., Taira, M. and Osada, S. (2009) Progranulin, a secreted tumorigenesis and dementia-related factor, regulates mouse hair growth. *J Dermatol Sci* 53(3): 234-6.

Hamada, S., Hara, K., Hamada, T., Yasuda, H., Moriyama, H., Nakayama, R., Nagata, M. and Yokono, K. (2009) Upregulation of the mammalian target of rapamycin complex 1 pathway by Ras homolog enriched in brain in pancreatic beta-cells leads to increased beta-cell mass and prevention of hyperglycemia. *Diabetes* 58(6): 1321-32.

Fukatsu, Y., Noguchi, T., Hosooka, T., Ogura, T., Kotani, K., Abe, T., Shibakusa, T., Inoue, K., Sakai, M., Tobimatsu, K., Inagaki, K., Yoshioka, T., Matsuo, M., Nakae, J., Matsuki, Y., Hiramatsu, R., Kaku, K., Okamura, H., Fushiki, T. and Kasuga, M. (2009) Muscle-specific overexpression of heparin-binding epidermal growth factor-like growth factor increases peripheral glucose disposal and insulin sensitivity. *Endocrinology* 150(6): 2683-91.

Imai, T., Yamazaki, T., Kobayakawa, R., Kobayakawa, K., Abe, T., Suzuki, M. and Sakano, H. (2009) Pre-target axon sorting establishes the neural map topography. *Science* 325(5940): 585-90.

Islam, S. M., Shinmyo, Y., Okafuji, T., Su, Y., Naser, I. B., Ahmed, G., Zhang, S., Chen, S., Ohta, K., Kiyonari, H., Abe, T., Tanaka, S., Nishinakamura, R., Terashima, T., Kitamura, T. and Tanaka, H. (2009) Draxin, a repulsive guidance protein for spinal cord and forebrain commissures. *Science* 323(5912): 388-93.

Oda, H., Fujimoto, M., Patrick, M. S., Chida, D., Sato, Y., Azuma, Y., Aoki, H., Abe, T., Suzuki, H. and Shirai, M. (2009) RhoH plays critical roles in Fc epsilon RI-dependent signal transduction in mast cells. *J Immunol* 182(2): 957-62.

Patrick, M. S., Oda, H., Hayakawa, K., Sato, Y., Eshima, K., Kirikae, T., Iemura, S., Shirai, M., Abe, T., Natsume, T., Sasazuki, T. and Suzuki, H. (2009) Gasp, a Grb2-associating protein, is critical for positive selection of thymocytes. *Proc Natl Acad Sci U S A* 106(38): 16345-50.

Tao, H., Suzuki, M., Kiyonari, H., Abe, T., Sasaoka, T. and Ueno, N. (2009) Mouse *prickle1*, the homolog of a PCP gene, is essential for epiblast apical-basal polarity. *Proc Natl Acad Sci U S A* 106(34): 14426-31.

Watanabe, N., Hiramatsu, K., Miyamoto, R., Yasuda, K., Suzuki, N., Oshima, N., Kiyonari, H., Shiba, D., Nishio, S., Mochizuki, T., Yokoyama, T., Maruyama, S., Matsuo, S., Wakamatsu, Y. and Hashimoto, H. (2009) A murine model of neonatal diabetes mellitus in *Glis3*-deficient mice. *FEBS Lett* 583(12): 2108-13.

Ohno, M., Hiraoka, Y., Matsuoka, T., Tomimoto, H., Takao, K., Miyakawa, T., Oshima, N., Kiyonari, H., Kimura, T., Kita, T. and Nishi, E. (2009) Nardilysin regulates axonal maturation and myelination in the central and peripheral nervous system. *Nat Neurosci* 12(12): 1506-13.

Noguchi, Y., Hirabayashi, T., Katori, S., Kawamura, Y., Sanbo, M., Hirabayashi, M.,



Kiyonari, H., Nakao, K., Uchimura, A. and Yagi, T. (2009) Total expression and dual gene-regulatory mechanisms maintained in deletions and duplications of the Pcdha cluster. *J Biol Chem* 284(46): 32002-14.

Nishioka, N., Inoue, K., Adachi, K., Kiyonari, H., Ota, M., Ralston, A., Yabuta, N., Hirahara, S., Stephenson, R. O., Ogonuki, N., Makita, R., Kurihara, H., Morin-Kensicki, E.M., Nojima, H., Rossant, J., Nakao, K., Niwa, H. and Sasaki, H. (2009) The Hippo signaling pathway components Lats and Yap pattern Tead4 activity to distinguish mouse trophectoderm from inner cell mass. *Dev Cell* 16(3): 398-410.

Nishide, K., Nakatani, Y., Kiyonari, H. and Kondo, T. (2009) Glioblastoma formation from cell population depleted of Prominin1-expressing cells. *PLoS One* 4(8): e6869.

Imuta, Y., Nishioka, N., Kiyonari, H. and Sasaki, H. (2009) Short limbs, cleft palate, and delayed formation of flat proliferative chondrocytes in mice with targeted disruption of a putative protein kinase gene, Pkdcc (AW548124). *Dev Dyn* 238(1): 210-22.

## 2008

Nishino, N., Tamori, Y., Tateya, S., Kawaguchi, T., Shibakusa, T., Mizunoya, W., Inoue, K., Kitazawa, R., Kitazawa, S., Matsuki, Y., Hiramatsu, R., Masubuchi, S., Omachi, A., Kimura, K., Saito, M., Amo, T., Ohta, S., Yamaguchi, T., Osumi, T., Cheng, J., Fujimoto, T., Nakao, H., Nakao, K., Aiba, A., Okamura, H., Fushiki, T. and Kasuga, M. (2008) FSP27 contributes to efficient energy storage in murine white adipocytes by promoting the formation of unilocular lipid droplets. *J Clin Invest* 118(8): 2808-21.

Nakamura, K., Ichise, H., Nakao, K., Hatta, T., Otani, H., Sakagami, H., Kondo, H. and Katsuki, M. (2008) Partial functional overlap of the three ras genes in mouse embryonic development. *Oncogene* 27(21): 2961-8.

Yamanaka, Y., Heike, T., Kumada, T., Shibata, M., Takaoka, Y., Kitano, A., Shiraishi, K., Kato, T., Nagato, M., Okawa, K., Furushima, K., Nakao, K., Nakamura, Y., Taketo, M.M., Aizawa, S. and Nakahata, T. (2008) Loss of Borealin/DasraB leads to defective cell proliferation, p53 accumulation and early embryonic lethality. *Mech Dev.* 125(5-6): 441-50.

Konno, D., Shioi, G., Shitamukai, A., Mori, A., Kiyonari, H., Miyata, T. and Matsuzaki, F. (2008) Neuroepithelial progenitors undergo LGN-dependent planar divisions to maintain self-renewability during mammalian neurogenesis. *Nat Cell Biol* 10(1): 93-101.

Yano, M., Kuroda, N., Han, H., Meguro-Horike, M., Nishikawa, Y., Kiyonari, H., Maemura, K., Yanagawa, Y., Obata, K., Takahashi, S., Ikawa, T., Satoh, R., Kawamoto, H., Mouri, Y. and Matsumoto, M. (2008) Aire controls the differentiation program of thymic epithelial cells in the medulla for the establishment of self-tolerance. *J Exp Med* 205(12): 2827-38.

Tamai, K., Toyoshima, M., Tanaka, N., Yamamoto, N., Owada, Y., Kiyonari, H., Murata, K., Ueno, Y., Ono, M., Shimosegawa, T., Yaegashi, N., Watanabe, M. and

Sugamura, K. (2008) Loss of hrs in the central nervous system causes accumulation of ubiquitinated proteins and neurodegeneration. *Am J Pathol* 173(6): 1806-17.

Takasato, M., Kobayashi, C., Okabayashi, K., Kiyonari, H., Oshima, N., Asashima, M. and Nishinakamura, R. (2008) Trb2, a mouse homolog of tribbles, is dispensable for kidney and mouse development. *Biochem Biophys Res Commun* 373(4): 648-52.

Suzuki, A., Sekiya, S., Onishi, M., Oshima, N., Kiyonari, H., Nakauchi, H. and Taniguchi, H. (2008) Flow cytometric isolation and clonal identification of self-renewing bipotent hepatic progenitor cells in adult mouse liver. *Hepatology* 48(6): 1964-78.

Sawada, A., Kiyonari, H., Ukita, K., Nishioka, N., Imuta, Y. and Sasaki, H. (2008) Redundant roles of Tead1 and Tead2 in notochord development and the regulation of cell proliferation and survival. *Mol Cell Biol* 28(10): 3177-89.

Nishioka, N., Yamamoto, S., Kiyonari, H., Sato, H., Sawada, A., Ota, M., Nakao, K. and Sasaki, H. (2008) 'Tead4 is required for specification of trophectoderm in pre-implantation mouse embryos', *Mech Dev* 125(3-4): 270-83.

Nakae, J., Cao, Y., Oki, M., Orba, Y., Sawa, H., Kiyonari, H., Iskandar, K., Suga, K., Lombes, M. and Hayashi, Y. (2008) Forkhead transcription factor FoxO1 in adipose tissue regulates energy storage and expenditure. *Diabetes* 57(3): 563-76.

Matsumoto, M., Zhou, Y., Matsuo, S., Nakanishi, H., Hirose, K., Oura, H., Arase, S., Ishida-Yamamoto, A., Bando, Y., Izumi, K., Kiyonari, H., Oshima, N., Nakayama, R., Matsushima, A., Hirota, F., Mouri, Y., Kuroda, N., Sano, S. and Chaplin, D.D. (2008) Targeted deletion of the murine corneodesmosin gene delineates its essential role in skin and hair physiology. *Proc Natl Acad Sci U S A* 105(18): 6720-4.

Kousaka, K., Kiyonari, H., Oshima, N., Nagafuchi, A., Shima, Y., Chisaka, O. and Uemura, T. (2008) Slingshot-3 dephosphorylates ADF/cofilin but is dispensable for mouse development. *Genesis* 46(5): 246-55.

Aoki, M., Kiyonari, H., Nakamura, H. and Okamoto, H. (2008) R-spondin2 expression in the apical ectodermal ridge is essential for outgrowth and patterning in mouse limb development. *Dev Growth Differ* 50(2): 85-95.

Seino, Y., Miki, T., Kiyonari, H., Abe, T., Fujimoto, W., Kimura, K., Takeuchi, A., Takahashi, Y., Oiso, Y., Iwanaga, T. and Seino, S. (2008) Isx participates in the maintenance of vitamin A metabolism by regulation of beta-carotene 15,15'-monooxygenase (Bcmo1) expression. *J Biol Chem* 283(8): 4905-11.

Shigetani, Y., Howard, S., Guidato, S., Furushima, K., Abe, T. and Itasaki, N. (2008) Wise promotes coalescence of cells of neural crest and placode origins in the trigeminal region during head development. *Dev Biol* 319(2): 346-58.

## 2007

Furushima, K., Yamamoto, A., Nagano, T., Shibata, M., Miyachi, H., Abe, T., Ohshima, N., Kiyonari, H. and Aizawa, S. (2007) Mouse homologues of Shisa antagonistic to Wnt and Fgf signalings. *Dev Biol* 306(2): 480-92.

Kurotaki, Y., Hatta, K., Nakao, K., Nabeshima, Y. and Fujimori, T. (2007) Blastocyst Axis Is Specified Independently of Early Cell Lineage But Aligns with the ZP Shape. *Science* 316(5825): 719-23.

Ogawa, M., Miyakawa, T., Nakamura, K., Kitano, J., Furushima, K., Kiyonari, H., Nakayama, R., Nakao, K., Moriyoshi, K. and Nakanishi, S. (2007) Altered sensitivities to morphine and cocaine in scaffold protein tamalin knockout mice. *Proc Natl Acad Sci U S A* 104(37): 14789-94.

Fujishima, K., Kiyonari, H., Kurisu, J., Hirano, T. and Kengaku, M. (2007) Targeted disruption of Sept3, a heteromeric assembly partner of Sept5 and Sept7 in axons, has no effect on developing CNS neurons. *J Neurochem* 102(1): 77-92.

Chan, T., Kondow, A., Hosoya, A., Hitachi, K., Yukita, A., Okabayashi, K., Nakamura, H., Ozawa, H., Kiyonari, H., Michiue, T., Ito, Y. and Asashima, M. (2007) Ripply2 is essential for precise somite formation during mouse early development. *FEBS Lett* 581(14): 2691-6.

Sugiyama, D., Ogawa, M., Nakao, K., Osumi, N., Nishikawa, S., Nishikawa, S., Arai, K., Nakahata, T. and Tsuji, K. (2007) B cell potential can be obtained from pre-circulatory yolk sac, but with low frequency. *Dev Biol* 301(1): 53-61.

Yana, I., Sagara, H., Takaki, S., Takatsu, K., Nakamura, K., Nakao, K., Katsuki, M., Taniguchi, S., Aoki, T., Sato, H., Weiss, S.J., Seiki, M. (2007) Crosstalk between neovessels and mural cells directs the site-specific expression of MT1-MMP to endothelial tip cells. *J Cell Sci* 120: 1607-1614.

Aiba, A., Inokuchi, K., Ishida, Y., Itohara, S., Kobayashi, K., Masu, M., Mishina, M., Miyakawa, T., Mori, H., Nakao, K., Obata, Y., Sakimura, K., Shiroishi, T., Wada, K. and Yagi, T. (2007) Mouse liaison for integrative brain research. *Neurosci Res* 58(2): 103-4.

Wei, P., Satoh, T., Edamatsu, H., Aiba, A., Setsu, T., Terashima, T., Kitazawa, S., Nakao, K., Yoshikawa, Y., Tamada, M. and Kataoka, T. (2007) Defective vascular morphogenesis and mid-gestation embryonic death in mice lacking RA-GEF-1. *Biochem Biophys Res Commun* 363(1): 106-12.

Sakata, D., Taniguchi, H., Yasuda, S., Adachi-Morishima, A., Hamazaki, Y., Nakayama, R., Miki, T., Minato, N. and Narumiya, S. (2007) Impaired T lymphocyte trafficking in mice deficient in an actin-nucleating protein, mDia1. *J Exp Med* 204(9): 2031-8.

Takasaki, N., Kurokawa, D., Nakayama, R., Nakayama, J. and Aizawa, S. (2007) Acetylated YY1 regulates Otx2 expression in anterior neuroectoderm at two cis-sites 90 kb apart. *EMBO J* 26(6): 1649-59.

Takashima, Y., Era, T., Nakao, K., Kondo, S., Kasuga, M., G. Smith, Austin. and Nishikawa, S. (2007) Neuroepithelial Cells Supply an Initial Transient Wave of MSC Differentiation. *Cell* 129(7): 1377-88.

Niwa, Y., Masamizu, Y., Liu, T., Nakayama, R., Deng, Chu-Xia. and Kageyama, R. (2007) The Initiation and Propagation of Hes7 Oscillation Are Cooperatively Regulated by Fgf and Notch Signaling in the Somite Segmentation Clock. *Dev Cell*. 13(2): 298-304.

Nakao, H., Nakao, K., Kano, M. and Aiba, A. (2007) Metabotropic glutamate receptor subtype-1 is essential for motor coordination in the adult cerebellum. *Neurosci Res* 57(4): 538-43.

Yoshikawa, Y., Satoh, T., Tamura, T., Wei, P., Bilasy, S E., Edamatsu, H., Aiba, A., Katagiri, K., Kinashi T., Nakao, K. and Kataoka, T. (2007) The

M-Ras-RA-GEF-2-Rap1 Pathway Mediates Tumor Necrosis Factor- $\alpha$ -dependent Regulation of Integrin Activation in Splenocytes. *Mol Bio Cell* 18(8): 2949-59.

Ohtani, N., Imamura, Y., Yamakoshi, K., Hirota, F., Nakayama, R., Kubo, Y., Ishimaru, N., Takahashi, A., Hirao, A., Shimizu, T., David J. Mann, Saya, H., Hayashi, Y., Arase, S., Matsumoto, M., Nakao, K., and Hara, E. (2007) Visualizing the dynamics of p21Waf1/Cip1 cyclin-dependent kinase inhibitor expression in living animals. *Proc Natl Acad Sci USA* 104(38): 15034-9.

## 2006

Kurokawa, D., Sakurai, Y., Inoue, A., Nakayama, R., Takasaki, N., Suda, Y., Miyake, T., Chris T. Amemiya. and Aizawa, S. (2006) Evolutionary constraint on Otx2 neuroectoderm enhancers-deep conservation from skate to mouse and unique divergence in teleost. *Proc Natl Acad Sci USA* 103(51): 19350-5.

Hirata, T., Nakazawa, M., Yoshihara, S., Miyachi, H., Kitamura, K., Yoshihara, Y. and Hibi, M. (2006) Zinc-finger gene Fez in the olfactory sensory neurons regulates development of the olfactory bulb non-cell-autonomously. *Development* 133(8): 1433-43.

Hirata, T., Nakazawa, M., Muraoka, O., Nakayama, R., Suda, Y. and Hibi, M. (2006) Zinc-finger genes Fez and Fez-like function in the establishment of diencephalon subdivisions. *Development* 133: 3993-4004.

Oda, M., Yamagiwa, A., Yamamoto, S., Nakayama, T., Tsumura, A., Sasaki, H., Nakao, K., Li, E. and Okano, M. (2006) DNA methylation regulates long-range gene silencing of an X-linked homeobox gene cluster in a lineage-specific manner. *Genes Dev* 20(24): 3382-94.

Sakaguchi, T., Nishimoto, M., Miyagi, S., Iwama, A., Morita, Y., Iwamori, N., Nakauchi, H., Kiyonari, H., Muramatsu, M. and Okuda, A. (2006) Putative "stemness" gene jam-B is not required for maintenance of stem cell state in embryonic, neural, or hematopoietic stem cells. *Mol Cell Biol* 26(17): 6557-70.

Mazaki, Y., Hashimoto, S., Tsujimura, T., Morishige, M., Hashimoto, A., Aritake, K., Yamada, A., Nam, J. M., Kiyonari, H., Nakao, K. and Sabe, H. (2006) Neutrophil direction sensing and superoxide production linked by the GTPase-activating protein GIT2. *Nat Immunol* 7(7): 724-31.

Ikeya, M., Kawada, M., Kiyonari, H., Sasai, N., Nakao, K., Furuta, Y. and Sasai, Y. (2006) Essential pro-Bmp roles of crossveinless 2 in mouse organogenesis.



Development 133(22): 4463-73.

Hirano, M., Kiyonari, H., Inoue, A., Furushima, K., Murata, T., Suda, Y. and Aizawa, S. (2006) A new serine/threonine protein kinase, Omphk1, essential to ventral body wall formation. *Dev Dyn* 235(8): 2229-37.

Hirai, S., Cui de, F., Miyata, T., Ogawa, M., Kiyonari, H., Suda, Y., Aizawa, S., Banba, Y. and Ohno, S. (2006) The c-Jun N-terminal kinase activator dual leucine zipper kinase regulates axon growth and neuronal migration in the developing cerebral cortex. *J Neurosci* 26(46): 11992-2002.

## 2005

Furushima, K., Murata, T., Kiyonari, H. and Aizawa, S. (2005) Characterization of Opr deficiency in mouse brain: subtle defects in dorsomedial telencephalon and medioventral forebrain. *Dev Dyn* 232(4): 1056-61.

Terada, K., Yomogida, K., Imai, T., Kiyonari, H., Takeda, N., Kadomatsu, T., Yano, M., Aizawa, S. and Mori, M. (2005) A type I DnaJ homolog, DjA1, regulates androgen receptor signaling and spermatogenesis. *EMBO J* 24(3): 611-22.

Nishioka, N., Nagano, S., Nakayama, R., Kiyonari, H., Ijiri, T., Taniguchi, K., Shawlot, W., Hayashizaki, Y., Westphal, H., Behringer, R. R., Matsuda, Y., Sakoda, S., Kondoh, H. and Sasaki, H. (2005) Ssdp1 regulates head morphogenesis of mouse embryos by activating the Lim1-Ldb1 complex. *Development* 132(11): 2535-46.

Ikeya, M., Kawada, M., Nakazawa, Y., Sakuragi, M., Sasai, N., Ueno, M., Kiyonari, H., Nakao, K. and Sasai, Y. (2005) Gene disruption/knock-in analysis of mONT3: vector construction by employing both in vivo and in vitro recombinations. *Int J Dev Biol* 49(7): 807-23.

Hirano, M., Murata, T., Furushima, K., Kiyonari, H., Nakamura, M., Suda, Y. and Aizawa, S. (2005) cfm is a novel gene uniquely expressed in developing forebrain and midbrain, but its null mutant exhibits no obvious phenotype. *Gene Expr Patterns* 5(3): 439-44.

Kimura, J., Suda, Y., Kurokawa, D., Houssain, ZM., Nakamura, M., Takahashi, M., Hara, A. and Aizawa, S. (2005) Emx2 and Pax6 Function in Cooperation with Otx2 and Otx1 to Develop Caudal Forebrain Primordium That Includes Future Archipallium. *J Neurosci* 25(21): 5097-108.

Tomemori, Y., Ichiba, M., Kusumoto, A., Mizuno, E., Sato, D., Muroya, S., Nakamura, M., Kawaguchi, H., Yoshida, H., Ueno, S., Nakao, K., Nakamura, K., Aiba, A., Katsuki, M. and Sano, A. (2005) A gene-targeted mouse model for chorea-acanthocytosis. *J*

Neurochem 92(4): 759-66.

Kassai, H., Aiba, A., Nakao, K., Nakamura, K., Katsuki, M., Xiong, WH., Yau, KW., Imai, H., Shichida, Y., Satomi, Y., Takao, T., Okano, T. and Fukada, Y. (2005) Farnesylation of Retinal Transducin Underlies Its Translocation during Light Adaptation. *Neuron* 47(4): 529-39.

Hara, T., Saito, Y., Hirai, T., Nakamura, K., Nakao, K., Katsuki, M. and Chida, K. (2005) Deficiency of protein kinase Calpha in mice results in impairment of epidermal hyperplasia and enhancement of tumor formation in two-stage skin carcinogenesis. *Cancer Res.* 65(16): 7356-62.

Hiratsuka, S., Nakao, K., Nakamura, K., Katsuki, M., Maru, Y. and Shibuya, M. (2005) Membrane fixation of vascular endothelial growth factor receptor 1 ligand-binding domain is important for vasculogenesis and angiogenesis in mice. *Mol Cell Biol* 25(1): 346-54.

Hiratsuka, S., Kataoka, Y., Nakao, K., Nakamura, K., Morikawa, S., Tanaka, S., Katsuki, M., Maru, Y. and Shibuya, M. (2005) Vascular endothelial growth factor A (VEGF-A) is involved in guidance of VEGF receptor-positive cells to the anterior portion of early embryos. *Mol Cell Biol* 25(1): 355-63.

Sawada, A., Nishizaki, Y., Sato, H., Yada, Y., Nakayama, R., Yamamoto, S., Nishioka, N., Kondoh, H. and Sasaki H. (2005) Tead proteins activate the *Foxa2* enhancer in the node in cooperation with a second factor. *Development* 132(21): 4719-29.

Kimura-Yoshida, C., Nakano, H., Okamura, D., Nakao, K., Yonemura, S., Belo, J.A., Aizawa, S., Matsui, Y. and Matsuo, I. (2005) Canonical Wnt signaling and its antagonist regulate anterior-posterior axis polarization by guiding cell migration in mouse visceral endoderm. *Dev Cell* 9(5): 639-50.

Saito, H., Komada, M., Suzuki, M., Nakayama, R., Motoyama, J., Shiota, K. and Ishibashi, M. (2005) Expression of the mouse *Fgf15* gene is directly initiated by Sonic

hedgehog signaling in the diencephalon and midbrain. *Dev Dyn* 232(2): 282-92.

Tada, S., Era, T., Furusawa, C., Sakurai, H., Nishikawa, S., Kinoshita, M., Nakao, K., Chiba, T. and Nishikawa, S. (2005) Characterization of mesendoderm: a diverging point of the definitive endoderm and mesoderm in embryonic stem cell differentiation culture. *Development* 132(19): 4363-74.

## 2004

Okita, K., Kiyonari, H., Nobuhisa, I., Kimura, N., Aizawa, S. and Taga, T. (2004) Targeted disruption of the mouse ELYS gene results in embryonic death at peri-implantation development. *Genes Cells* 9(11): 1083-91.

Murata, T., Furushima, K., Hirano, M., Kiyonari, H., Nakamura, M., Suda, Y. and Aizawa, S. (2004) ang is a novel gene expressed in early neuroectoderm, but its null mutant exhibits no obvious phenotype. *Gene Expr Patterns* 5(2): 171-8.

Kurokawa, D., Takasaki, N., Kiyonari, H., Nakayama, R., Kimura-Yoshida, C., Matsuo, I. and Aizawa, S. (2004) Regulation of Otx2 expression and its functions in mouse epiblast and anterior neuroectoderm. *Development* 131(14): 3307-17.

Kurokawa, D., Kiyonari, H., Nakayama, R., Kimura-Yoshida, C., Matsuo, I. and Aizawa, S. (2004) Regulation of Otx2 expression and its functions in mouse forebrain and midbrain. *Development* 131(14): 3319-31.

Sato, M., Tabata, T., Hashimoto, K., Nakamura, K., Nakao, K., Katsuki, M., Kitano, J., Moriyoshi, K., Kano, M. and Nakanishi, S. (2004) Altered agonist sensitivity and desensitization of neuronal mGluR1 responses in knock-in mice by a single amino acid substitution at the PKC phosphorylation site. *Eur J Neurosci* 20(4): 947-55.

Hirata, T., Suda, Y., Nakao, K., Narimatsu, M., Hirano, T. and Hibi, M. (2004) Zinc finger gene fez-like functions in the formation of subplate neurons and thalamocortical axons. *Dev Dyn* 230(3): 546-56.

## 2003

Terada, K., Ueno, S., Yomogida, K., Imai, T., Kiyonari, H., Takeda, N., Yano, M., Abe, S., Aizawa, S. and Mori, M. (2003) Expression of Tom34 splicing isoforms in mouse testis and knockout of Tom34 in mice. *J Biochem* 133(5): 625-31.

Chida, K., Hara, T., Hirai, T., Konishi, C., Nakamura, K., Nakao, K., Aiba, A., Katsuki, M. and Kuroki, T. (2003) Disruption of protein kinase Ceta results in impairment of wound healing and enhancement of tumor formation in mouse skin carcinogenesis. *Cancer Res* 63(10): 2404-8.

Takaki, S., Tezuka, Y., Sauer, K., Kubo, C., Kwon, SM., Armstead, E., Nakao, K., Katsuki, M., Perlmutter, RM. and Takatsu, K. (2003) Impaired lymphopoiesis and altered B cell subpopulations in mice overexpressing Lnk adaptor protein. *J Immunol* 170(2): 703-10.